Draft version February 2, 2008
Preprint typeset using LaTeX style emulateapj v. 9/08/03

arXiv:astro-ph/0312550v1 19 Dec 2003

A DISK GALAXY OF OLD STARS AT Z ~ 2.5[1]

Alan Stockton
Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

Gabriela Canalizo
Institute of Geophysics and Planetary Physics and Department of Earth Sciences, University of California, Riverside, CA 95521

Toshinori Maihara
Department of Astronomy, Kyoto University, Kitashirakawa-Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

Draft version February 2, 2008

ABSTRACT

We describe observations of a galaxy in the field of the $z = 2.483$ radio galaxy 4C 23.56, photometrically selected to have a spectral-energy distribution consistent with an old stellar population at the redshift of the radio galaxy. Exploration of redshift—stellar-population—reddening constraints from the photometry indicates that the galaxy is indeed at a redshift close to that of 4C 23.56, that the age of the most recent significant star formation is roughly $\gtrsim 2$ Gyr, and that reddening is fairly modest, with more reddening required for the younger end of stellar age range. From analysis of a deep adaptive-optics image of the galaxy, we find that an $r^{1/4}$-law profile, common for local spheroidal galaxies, can be excluded quite strongly. On the other hand, a pure exponential profile fits remarkably well, while the best fit is given by a Sérsic profile with index $n = 1.49$. Reconstruction of the two-dimensional form of the galaxy from the best-fit model is consistent with a disk galaxy with neither a significant bulge component nor gross azimuthal structure. The assembly of roughly $2L^*$ of old stars into such a configuration this early in the history of the universe is not easily explainable by any of the currently popular scenarios for galaxy formation. A galaxy with these properties would seem to require smooth but rapid infall of the large mass of gas involved, followed by a burst of extremely vigorous and efficient star formation in the resulting disk.

Subject headings: galaxies: high-redshift—galaxies: formation—galaxies: evolution

1. INTRODUCTION

Our understanding of the mechanisms and timescales of the formation of the very first luminous galaxies in the universe is still very uncertain. Many detailed studies of local luminous early-type galaxies favor formation at very early epochs and within a very short time span (*e.g.,* Peebles 2002 and references therein). In particular, the α elements are enhanced in luminous ellipticals relative to Fe (*e.g.,* Worthy, Faber, & González 1992), this enhancement is strongly correlated with the central velocity dispersion σ (Trager et al. 2000), and the relation is similar for field and cluster ellipticals (Bernardi et al. 1998; Colless et al. 1999; Thomas, Maraston, & Bender 2002). Galaxies with the oldest stellar populations are both the most luminous and the most α enhanced (Thomas, Maraston, & Bender 2002). If these claims are valid, the most luminous galaxies formed within a quite short period very early in the history of the universe, less luminous early-type galaxies formed more gradually and somewhat later, and differences in star formation histories between ellipticals in the field and those in clusters are relatively minor. This last statement is also supported by fundamental-plane analyses (*e.g.,* van Dokkum & Ellis 2003; Treu et al. 2001; but see also Treu et al. 2002). Such conclusions appear to be difficult to reconcile with standard interpretations of the results of the semi-analytic cold-dark-matter (CDM) models (*e.g.,* Thomas & Kauffmann 1999; Kauffmann & Haehnelt 2000), which predict that, for the currently favored low-density cosmology ($H_0 \sim 70$, $\Omega_m \sim 0.3$, $\Omega_{vac} \sim 0.7$; assumed throughout this paper), most of the present-day bright ellipticals will have undergone final assembly between $z = 1$ and $z = 2$. However, there are at least two caveats to this statement: (1) One of the least certain parts of the semi-analytic CDM models is the treatment of star formation and resulting feedback effects, and it is possible that different assumptions in this area could mitigate at least some of the discrepancy; and (2) *assembly* at relatively low redshifts is not necessarily inconsistent with formation of most of the stars at much higher redshifts. Nevertheless, both the color-luminosity relationship for spheroids (Bower, Lucey, & Ellis 1992; Ellis et al. 1997; van Dokkum et al. 2000) and the correlation of α-element enhancement with luminosity show that the formation of the stellar population and the assembly of the galaxy cannot be completely independent processes. As Peacock (1999) has succinctly put it, "It seems as if the stars in ellipticals were formed at a time when the depth of the potential well that they would eventually inhabit was already determined."

---

[1] Based in part on data collected at Subaru Telescope, which is operated by the National Astronomical Observatory of Japan. Some of the data presented herein were obtained at the W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation.

In this context, it is of considerable interest to seek out and to investigate the galaxies with the oldest stellar populations at any given redshift. While such galaxies may not be typical, they can establish one end of the range of realistic formation scenarios, and they likely have less complicated star-formation histories than most. Furthermore, they may, in fact, be more representative of the formation of the bulk of the stars in early-type galaxies generally than might at first seem likely, since a miniscule contamination of young stars in a galaxy can drastically alter its rest-frame ultraviolet colors and eliminate such galaxies from color-selected samples.

There has been much recent activity in observations and discussion of so-called *extremely red objects* (EROs; *e.g.,* Liu et al. 2000; Daddi et al. 2000, 2003; Stockton 2001; Cimatti et al. 2003; Yan & Thompson 2003; Miyazaki et al. 2003; Gilbank et al. 2003; earlier references can be found in these papers). These objects, which are often defined to have $R-K \gtrsim 6$ or $I-K > 4$, include two disparate classes. Some are highly reddened high-redshift starbursts (*e.g.,* Dey et al. 1999). The remainder are almost exclusively galaxies with $z \gtrsim 1.2$ whose stellar populations are already ancient and have little admixture from more recent star formation (Stockton, Kellogg, & Ridgway 1995; Dunlop et al. 1996; Spinrad et al. 1997; Liu et al. 2000; Soifer et al. 1999; Stockton 2001). For convenience, we shall henceforth simply refer to this latter class as "old galaxies" (OGs); but note once again that this term refers to the age of the stellar population, with no necessary implication regarding the age of assembly of the galaxy itself, aside from the sorts of constraints mentioned above. In any case, these OGs seem to represent the earliest major episodes of star formation in the universe.

The actual ages of the dominant stellar population in the small number of identified OGs that have been studied in detail have generated some controversy because of disagreements about spectral synthesis models and their interpretation (see, *e.g.,* Dunlop 1999 and references therein). However, a minimum age of 2 Gyr for the oldest objects at $z \sim 1.5$ is accepted by virtually everyone, and there are good reasons for believing that some, at least, have ages between 3 and 4 Gyr (Dunlop 1999; Stockton 2001; Nolan, Dunlop, & Jimenez 2001; Nolan et al. 2002). If these latter age estimates are correct, then the precursors of these galaxies will still have essentially fully formed stellar populations at higher redshifts. For example, for our assumed cosmological parameters, galaxies at $z = 2.5$ will be ~ 1.6 Gyr younger than at $z = 1.5$. This means that $z = 1.5$ galaxies for which the last major episode of star formation occurred between 3 and 4 Gyr earlier might be expected to be fairly well settled and have small average internal extinction at $z = 2.5$, and probably at even higher redshifts.

Many surveys of EROs have made the assumption, explicit or not, that a dominant spheroidal component (in mass, if not in light) is a necessary condition for the presence of an old stellar population at high redshifts (*e.g.,* Moriando, Cimatti, & Daddi 2000; Smith et al. 2002b). In this paper, we describe the first object among those found in our search for OGs in radio source fields at $z \sim 2.5$ that we have been able to investigate in some detail, and we show that it is a luminous disk galaxy, with little or no bulge component, comprising very old stars.

2. SELECTING GALAXIES WITH OLD STELLAR POPULATIONS IN RADIO-SOURCE FIELDS AT $Z \sim 2.5$

We have recently concluded a search for OGs in the fields of quasars at $z \sim 1.5$ (Stockton 2001, Stockton et al., in preparation). Radio source fields are generally likely to include regions of higher-than-average density in the early universe (*e.g.,* Best et al. 2003), in which processes of galaxy formation and evolution are expected to have proceeded more rapidly than in the field. More importantly, looking for companions to radio sources at a specific redshift allows us to choose redshifts for which the photometric diagnostics from standard broadband filters give the cleanest separation between old galaxy populations and other possible contaminants, such as the dusty starbursts mentioned above. The relevant key redshifts for our present purposes are $z \sim 1.5$ (4000 Å break just shortward of the J band) and $z \sim 2.5$ (break between J and H bands).

Assume that, for the brighter OGs from our sample at $z \sim 1.5$, essentially all of their stars formed within 500 Myr of the Big Bang (corresponding to a stellar population age of 3.7 Gyr at $z = 1.5$). For passive evolution of the stellar population, one finds from solar metallicity models (see Bruzual & Charlot 1993, 2003; we have used the 2000 versions of the models for analysis here unless otherwise specified) that the precursors of such galaxies at $z \sim 2.5$ would have $J \sim 22.4$, $H \sim 20.5$, and $K' \sim 19.5$. This assumption is deliberately an extreme one: if our assumed age (corresponding to $z_{formation} \sim 9$) is too high, then these magnitudes are all a bit pessimistic. Our approach has been first to obtain imaging to $K' \sim 20$ (10σ) and $J \sim 23$ (5σ), looking for objects in the field with $J-K' \sim 3$. If any likely candidates are present, we obtain deeper imaging at J and K', and include imaging in the H band. Most of the infrared observations (including all of those relevant to this paper) have been obtained with the infrared camera CISCO (Motohara et al. 2002) on the 8.2 m Subaru telescope on Mauna Kea. The detector is a Rockwell 1024×1024 Hg-Cd-Te Hawaii-2 array, with a pixel scale of $0\farcs105$ and thus a field of just under $2'$. Additional R-band imaging was obtained with ESI on the Keck II telescope.

Our sample comprises 87 radio-source fields from the Texas catalog (Douglas et al. 1996) selected with the help of the NASA/IPAC Extragalactic Database (NED) to have redshifts in the range $2.3 < z < 2.7$ and $A_B < 1$, based on the Galactic extinction estimates of Schlegel, Finkbeiner, & Davis (1998). The coverage we have been able to achieve so far on these fields has been largely determined by telescope scheduling and weather. We currently have adequate observations for 10 fields, of which at least 3 have objects that have well characterized spectral-energy distributions (SEDs) that closely match those expected for old stellar populations at the redshifts of the radio sources, with little intrinsic reddening. General results for this survey will be given elsewhere; here we concentrate on one galaxy, found in the field of the radio galaxy 4C 23.56, for which we have both strong constraints on the SED and adaptive-optics (AO) imaging. A finding chart for the object is shown in Fig. 1. Al-



Fig. 1.— Field of the radio galaxy 4C 23.56. The image was obtained with the CISCO IR camera on the Subaru telescope in the K' band. The galaxy with a SED indicating an old stellar population is indicated as KC68. The bright star to the lower-right side of the image was used as the AO guide star.



Fig. 2.— Spectral-energy distribution in the observed frame for 4C 23.56 KC68. The photometric bands are R, I, J, H, and K'; the I-band upper limit is from Knopp & Chambers (1996). The vertical bars show 1σ errors, and the horizontal bars indicate the FWHM of the filter. The upper limits shown are 3σ. All of the photometric points and upper limits have been corrected for Galactic extinction using the estimate of Schlegel, Finkbeiner, & Davis (1998). The colored traces are examples of best-fit redshifts, stellar populations, and extinctions, as discussed in the text.

though it was found independently in our survey, this galaxy was previously noted as a very red galaxy by Knopp & Chambers (1996) in their comprehensive study of the field of 4C 23.56. The galaxy is number 68 in their Table 4, and we will refer to it as 4C 23.56 KC68, or, more briefly, simply as KC68.

The photometry for 4C 23.56 KC68 is shown in Fig. 2. The magnitudes, prior to Galactic reddening correction, are $R > 26.7$ (3σ), $I > 26.5$ (3σ), $J = 23.38 \pm 0.12$, $H = 20.91 \pm 0.06$, $K' = 19.90 \pm 0.03$. We have used the photometric redshift code Hyperz[2] (Bolzonella, Miralles, & Pelló 2000) to explore possible fits to the photometric data, and some of these model SEDs are also shown in Fig. 2. The fitting procedure allowed redshifts $0 \leq z \leq 6$, internal extinction corrections $0 \leq A_V \leq 6$ (assuming a Calzetti et al. 2000 reddening law), and a full range of solar-metallicity stellar population models, including both single-burst models as well as models incorporating various rates of continuing star formation. Within this parameter space, several general conclusions emerged:



Fig. 3.— Run of χ^2 with z for model fits to the photometry for 4C 23.56 KC68. Models with different ranges of internal extinction have been separated. At each redshift, the χ^2 statistic for the best-fitting model has been plotted, so the ages and internal extinction vary along each trace.

1. Only single-burst models produced good fits. This is presumably because significant continuing star formation would weaken the sharp inflection shortward of the H-band point and/or violate the R and I-band upper limits.

2. The redshift is strongly constrained to the range $2.1 \leq z \leq 2.7$. Figure 3 shows the χ^2 statistic for the fits as a function of redshift.

3. When redshift was left as a free parameter, the fits tended to home in on ~ 2.35, with a 99% confidence interval $2.1 \lesssim z \lesssim 2.7$. If one assumes the galaxy to be a member of a cluster associated with 4C 23.56 and therefore fixes the redshift at 2.483, a fairly respectable solution exists with an age of 2.6 Gyr and a modest internal extinction $A_V = 0.16$ mag.

4. Although the lowest reduced χ^2 was obtained for an old (3.0 Gyr) population with $A_V \sim 0.14$, to some extent it is possible to trade younger populations for increased reddening. However, this exchange cannot be carried too far: for ages $\lesssim 1$ Gyr the rapid rise in the flux at rest-frame 2000 Å, coupled with the bluer colors longward of 4000 Å, prevents the underlying continuum from being

---

[2] `http://webast.ast.obs-mip.fr/hyperz/`

Table 1. Model Fits to Photometry of 4C 23.56 KC68

Model	z	Age[a] (Gyr)	A_V (mag)	χ^2
1	2.340	3.0	0.14	1.128
2	2.483	2.6	0.16	1.832
3	2.340	1.0	1.10	1.877

[a] Assuming solar metallicity. See text for further discussion.

massaged to give a reasonable fit to the photometry by any amount of Calzetti-like extinction.

Table 1 gives parameters for the three models shown in Fig. 2, which are representative of the range of plausible fits to the photometry.

Note that, because of the well-known age-metallicity degeneracy, we can obtain younger ages by allowing super-solar metallicities. Such a trade-off means, for example, that even though some of the ages in Table 1 exceed the age of the universe for the given redshift and our assumed cosmological parameters (*i.e.,* 2.75 Gyr at $z = 2.34$; 2.59 Gyr at $z = 2.48$), these excess ages can be considered proxies for high metallicities. At a redshift of 2.48, an L^* elliptical with a 2.1-Gyr-old solar-metallicity stellar population would have $K' \sim 20.5$ in the observed frame, so the observed $K' = 19.8$ (after correction for Galactic extinction) for KC68 implies a luminosity of about $2L^*$. This mass associated with this luminosity is not strongly dependent on metallicity, the differential effect between solar and twice solar metallicities being a reduction in the inferred mass by about 10%.

3. ADAPTIVE-OPTICS IMAGING OF 4C 23.56 KC68

The galaxy 4C 23.56 KC68 happens to lie $\sim 20''$ from a 14th mag star, and we have been able to obtain adaptive-optics (AO) imaging, using the Subaru 36-element, curvature-sensing AO system and the Infrared Camera and Spectrograph (IRCS; Kobayashi et al. 2000). The 1024×1024 Aladdin3 array had a pixel scale of $0''.0225$, giving a field of $23''.0$. The total exposure was 12,600 s, all in the K' filter. Figure 4 shows a section of the AO-imaged field. The image cores have FWHM $= 0''.10$, and there is no evidence for significant anisoplanicity effects across the relevant part of the field. Insets in Fig. 4 show larger-scale images of KC68 at two contrast levels. The galaxy clearly has a high degree of symmetry, showing that $\sim 2L^*$ galaxies, comprising relaxed old stellar populations, can be in place by the time the universe is only 2.6 Gyr old.

We can explore this morphology in more detail by making model fits to the image. We have used C. Y. Peng's GALFIT[3] routine (Peng et al. 2002) to fit two-dimensional models corresponding to de Vaucouleurs ($r^{1/4}$-law), exponential, and general Sérsic profiles(Sérsic 1968; Caon, Capaccioli, & D'Onofrio 1993), all convolved with the PSF. Figure 5 shows these models and the residuals after subtraction from the observed image. Although a pure exponential model is an excellent fit, the formal best fit is for a Sérsic model with an index $n = 1.49$, an effective radius $r_e = 0''.22$ (= 1.8 kpc), and an axial ratio $b/a = 0.33$. These numbers tend to confirm one's visual impression that the galaxy is disk-like in morphology. While one cannot exclude the possibility of a small $r^{1/4}$-law bulge component, all of our attempts to model the galaxy as a two-component ($r^{1/4}$-law + exponential) system resulted in the $r^{1/4}$-law-component being given an extremely large effective radius (regardless of the initial parameter settings), essentially smearing it into the sky background.

For a more quantitative view, Fig. 6 shows the surface-brightness profile of KC68, along with seeing-convolved model fits. It is clear from the left panel that an $r^{1/4}$-law model cannot adequately fit the profile, and that an exponential model gives a much better fit. The right panel shows the best Sérsic-profile fit.

4. SUMMARY AND DISCUSSION

The identification of this galaxy and the other two (mentioned in §2) at $z \sim 2.5$ that we have found so far with strikingly dominant old stellar populations demonstrates that at least some massive galaxies have completed virtually all of their star formation at sufficiently early times that they are no longer heavily dust-enshrouded at this redshift. Of these, at least 4C 23.56 KC68 has also achieved a relaxed regular morphology. The surprise is that this relaxed configuration seems to be very close to a pure disk, rather than to a spheroid, or even to a spheroid + disk.

There is abundant evidence that the large ($\sim$ 40%) fraction of S0 galaxies seen in rich clusters at the present epoch drops dramatically at higher redshifts (Dressler et al. 1997; Fasano et al. 2000), with the slack being taken up by spirals, and with the proportion of elliptical galaxies remaining roughly constant. The mechanisms for the apparent conversion of spirals into S0 galaxies have been widely discussed ever since Butcher & Oemler (1978) first noticed the increasing fraction of blue galaxies in clusters at high redshifts (see, *e.g.,* Bicker, Fritze-v. Alvensleben, & Fricke 2002 for a recent example).

However, there is also evidence to support the classical view that at least a significant fraction of the S0 population consists of galaxies that formed coevally with the oldest ellipticals (Tinsley & Gunn 1976; Ellis et al. 1997). Iye et al. (2000, 2003) have described an S0-like galaxy at $z \sim 1.5$. In addition, recent large surveys find that a substantial fraction of objects identified as EROs have disks (Smith et al. 2002a; Yan & Thompson 2003; Gilbank et al. 2003; see also Smith et al. 2002b), and the only galaxy in our own sample of galaxies at $z \sim 1.5$ with old stellar populations (Stockton 2001) for which we have good AO imaging also is dominated by an exponential profile (Stockton et al., in preparation). At a higher redshifts, Labbé et al. (2003) have identified a sample of large, disk-like galaxies, one of which has fairly red colors (Franx et al. 2003) and is at $z = 2.94$.

All of these cases appear still to have significant (and usually dominant) bulge components. Indeed, one of the principal points of Smith et al. (2002b) is that the galaxy they describe may be a spheroid in the process of acquiring a disk, validating a commonly proposed scenario for

---

[3] `http://zwicky.as.arizona.edu/∼cyp/work/galfit/galfit.html/`



FIG. 4.— Image of the field around 4C 23.56 KC68 obtained with the AO system and IRCS on the Subaru telescope. The main image has been smoothed with a Gaussian kernel with $\sigma = 2$ pixels. The insets show KC68 enlarged by a factor of 2: the lower-left inset shows the unsmoothed image at low contrast, and the lower-right inset shows an image at medium contrast, smoothed with a Gaussian kernel with $\sigma = 1$ pixel. A ghost image from the AO guide star is indicated.

the evolution of early-type spirals. In other words, the reason that this particular galaxy is classified as an ERO lies in the dominance of the bulge population and the weakness and low star-formation rate of the disk component. Many other galaxies classified as EROs also show discrete star-forming regions, but in these cases, too, the bulge population seems to dominate the colors used for the classification.

The galaxy we are discussing here poses a much more radical challenge to our understanding of the formation of old stellar populations at high redshifts: KC68 appears to be a $\sim 2L^*$ galaxy of old stars having a disk-like morphology, with no significant bulge component. As indicated below, this galaxy seems so difficult to fit into the standard cold-dark-matter (CDM) picture of early galaxy formation that it is worth briefly revisiting the steps by which we have reached this conclusion.

Could the redshift be seriously in error?—Certainly for low extinction, the photometric redshift seems quite unambiguous (see Fig. 3). For the Calzetti et al. (2000) reddening law, this conclusion remains for the highest extinctions we have considered. We cannot exclude the possibility that one could contrive a reddening law that could result in a reasonable fit at a different redshift, for sufficiently high extinction. However, the high degree of symmetry shown by the galaxy does not fit the fairly chaotic morphology usually shown by galaxies having large amounts of dust. We believe that a redshift outside of the range $2.1 \leq z \leq 2.7$ is extremely unlikely.

Are the stars really old?—We have seen in §2 that reddening alone cannot reproduce the observed SED, including the sharp inflection shortward of the H band and the tight constraints on the flux at rest-frame 2000 Å and 2400 Å given by the R and I-band upper limits. (This conclusion strictly assumes a Calzetti et al. 2000 reddening law; but the statement almost certainly holds for any plausible reddening law). It is necessary to have a strong break in the underlying spectrum, even if reddening is substantial. Younger ages produce progressively worse fits, and ages < 1 Gyr do not seem at all likely, provided only that the stellar initial mass function (IMF) is not pathologically deficient in high-mass stars. Of course, one can make the stellar population as young as one wishes by an *ad hoc* truncation of the IMF above about $1.8M_\odot$ (*e.g.,* Broadhurst & Bouwens 2000). For standard IMFs (*e.g.,* Salpeter or Miller-Scalo), increasing the metallicity above solar can potentially give a slightly younger age. Experiments with the recent Bruzual & Charlot (2003) models show that the gross SEDs of solar-metallicity models can be reproduced by models at 2.5 times solar metallicity with ages reduced by about 30%. However, any real stellar population will have a range of metallicities, and chemically consistent evolutionary synthesis models (*e.g.,* Fritze-v. Alvensleben 1999; Jimenez et al. 2000) show that populations with mass-weighted metallicities that are much higher than solar can still have integrated fluxes dominated by stars with sub-solar metallicities in the rest-frame near-UV, where line blanketing is important. This effect means that ages estimated from solar-



FIG. 5.— Model fits to 4C 23.56 KC68. The top-left image reproduces the observed image from Fig. 4. The three rows underneath show PSF-convolved de Vaucouleurs, exponential, and Sérsic models calculated with GALFIT (Peng et al. 2002). The first panel of each row shows the model, using the same stretch as for the observed image; the second panel shows the residual after subtraction of the model from the image, again at the same contrast; the third panel shows the residual after Gaussian smoothing with a $\sigma = 2$ pixel kernel and a factor of 3 contrast enhancement. Note that the de Vaucouleurs model systematically oversubtracts at both small and large radii while undersubtracting at intermediate radii. Differences between the exponential and Sérsic profiles are quite subtle, as expected from the small Sérsic index. The top-right panel shows the appearance of the Sérsic model before convolution with the PSF, which is presumably a good indication of the gross morphology of the galaxy.



Fig. 6.— The observed radial surface-brightness profile of 4C 23.56 KC68 is shown as the black points with error bars in both panels. In the left panel are plotted best-fit de Vaucouleurs (green) and exponential (red) profiles from two-dimensional models convolved with the PSF, along with the PSF itself (black dashed line). The exponential model is clearly a much better fit; the galaxy model itself (*i.e.,* before convolution) has an effective radius $r_e = 0.''12$ and an axial ratio $b/a = 0.35$. The right panel shows the best Sérsic profile fit, again after convolution with the PSF. The model has an index $n = 1.49$, an effective radius $r_e = 0.''22$, and an axial ratio $b/a = 0.33$.

metallicity models are unlikely to be serious overestimates (Dunlop 1999). In the case of KC68, there seems to be no way to avoid requiring almost all of the stars to be close to 1 Gyr old or older. Furthermore, such solutions are about a factor of 3 less probable than the best fit, which is for a very old (2–3 Gyr) population with little reddening. Thus, it is likely that the most recent significant star formation in 4C 23.56 KC68 occurred when the universe was < 1 Gyr old, although we cannot completely exclude somewhat later times. (Note that if we were to assume the fairly extreme solution of a 1-Gyr-old population with internal reddening of $A_V \sim 1.1$ mag, the star formation would have to be essentially complete by $z \sim 3.8$, when the age of the universe would have been 1.6 Gyr). The important point remains, in any case, that there cannot have been significant recent star formation.

Is the morphology really disk-like?—As we have stated in §3, we cannot exclude the possibility of a small $r^{1/4}$-law component, but any such component seems to be insignificant in terms of the total luminosity of the system. The small projected axial ratio and the good fit to a pure exponential profile indicate that the system is strongly dominated by a disk.

In sum, the most straightforward and plausible interpretation remains that 4C 23.56 KC68 is a well-settled disk of old stars at a redshift close to that of 4C 23.56 itself. If this interpretation is indeed correct, what are the consequences? Firstly, something very close to the classical picture of monolithic collapse must have occurred, with the important proviso that vigorous star formation did not begin until after the gas had settled into a disk. Secondly, star formation must have proceeded at an extremely rapid rate in order to form the required $\sim 3 \times 10^{11} M_\odot$ of stars over a time span that could not possibly be much more than 10^9 years and that was probably more like a few 10^8 years. Thirdly, the star formation and processes associated with it must have been very efficient at processing and expelling the gas in order to have strongly quenched any subsequent significant star formation.

In hierarchical formation models, including the standard CDM picture, galaxy morphologies are largely a reflection of their merger and accretion histories up to the epoch at which they are observed (*e.g.,* Steinmetz & Navarro 2003). Semi-analytic models of galaxy formation (*e.g.,* Kauffmann & Charlot 1998) suggest that most massive spheroids underwent final assembly between redshifts of 1 and 2, and that most large spiral disks were gradually acquired by smooth accretion of cold gas at redshifts ∼ 1 and lower, after the last major merger (Katz & Gunn 1991; Navarro & White 1994). There appears to be little place in this scenario for a galaxy like 4C 23.56 KC68, for at least two reasons: (1) the implied prodigious star formation rate, which has to have been sustained for a few 10^8 years *without* the help of a major merging event (as indicated by the lack of a detectable bulge), and (2) the building up of a massive galaxy apparently by smooth accretion of cold gas, without any evidence for significant hierarchical merging.

We of course cannot say from this one example whether similar galaxies are common at high redshifts. If they are, they would in one sense be the perfect progenitors for spheroidals: they comprise natural reservoirs of stars formed over very short periods of time in deep potential wells at early epochs. In short, their stars have all of the properties needed to explain the color-luminosity and alpha-element-luminosity relations for spheroidals, provided only that the morphological transformation from disks to spheroidals results from mergers of a small number of massive disk galaxies (Kauffmann & Charlot 1998). One obvious course for the future would be to explore the morphologies of galaxies selected in the same way as KC68 was with high resolution imaging, either with NICMOS on *HST*, or with laser-guide-star AO on large ground-based telescopes.

We thank Jenny Patience for assisting with some of the Subaru observations, and we thank the staff of the Subaru and Keck telescopes for their excellent support for our observing programs. We have benefited from discussions with Josh Barnes, John Kormendy, and Brent Tully. This research has been partially supported by NSF grant AST03-07335. It made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. The authors recognize the very significant cultural role that the summit of Mauna Kea has within the indigenous Hawaiian community and are grateful to have had the opportunity to conduct observations from it.

REFERENCES

Bernardi, M., et al. 1998, ApJ, 508, L143

Best, P. N., Lehnert, M. D., Miley, G. K., & Röttgering, H. J. A. 2003, MNRAS, 343, 1

Bicker, J., Fritze-v. Alvensleben, U., & Fricke, K. J. 2002, A&A, 387, 412

Bolzonella, M., Miralles, J.-M., & Pelló, R. 2000, A&A, 363, 476

Bower, R. G., Lucey, J. R., & Ellis, R. S. 1992, MNRAS, 254, 601

Broadhurst, T., & Bouwens, R. J. 2000, ApJ, 530, L53

Bruzual, G. & Charlot, S. 1993, ApJ, 405, 538

Bruzual, G. & Charlot, S. 2003, MNRAS, 344, 1000

Butcher, H., & Oemler, A. 1978, ApJ, 219, 18

Calzetti, D., Armus, L., Bohlin, R. C., Kinney, A. L., Kornneef, J., & Storchi-Bergmann, T. 2000, ApJ, 533, 682

Caon, N., Capaccioli, M., & D'Onofrio, M. 1993, MNRAS, 265, 1013

Cimatti, A., et al. 2003, A&A, in press

Colless, M., Burstein, D., Davies, R. L., McMahan, R. K., Saglia, R. P., & Wegner, G. 1999, MNRAS, 303, 813

Daddi, E., et al. 2000, A&A, 361, 535.

Daddi, E., Röttgering, H. J. A., Labbé, I., Rudnick, G., Franx, M., Moorwood, A. F. M., Rix, H. W., van der Werf, P. P., & van Dokkum, P. G. 2003, ApJ, 588, 50

Dey, A., et al. 1999, ApJ, 519, 610

Dickinson, M., Papovich, C., Ferguson, H. C., & Budavári, T. 2003, ApJ, 587, 25

Douglas, J. N., Bash, F. N., Bozyan, F. A., Torrence, G. W., & Wolfe, C. 1996, AJ, 111, 1945

Dressler, A., Oemler, Jr., A., Couch, W. J., Smail, I., Ellis, R. S., Barger, A., Butcher, H., Poggianti, B. M., & Sharples, R. M. 1997, ApJ, 490, 577

Dunlop, J. S. 1999, in The Hy-Redshift Universe, ASP Conf. Ser., 193, ed. A. J. Bunker & W. J. M. van Breugel (ASP, San Francisco), p. 133

Dunlop, J., Peacock, J., Spinrad, H., Dey, A., Jimenez, R., Stern, D., & Windhorst, R. 1996, Nature, 381, 581

Ellis, R. S., Smail, Ian, Dressler, A., Couch, W. J., Oemler, Jr., A. Butcher, H., & Sharples, R. M. 1997, ApJ, 483, 582

Fasano, G., Poggianti, B. M., Couch, W. J., Bettoni, D., Kjærgaard, P., & Moles, M. 2000, ApJ, 542, 673

Franx, M., et al. 2003, ApJ, 587, L79

Fritz-v. Alvensleben, U. 1999, in Spectrophotometric Dating of Stars and Galaxies, ASP Conf. Ser., 192, ed. I. Hubeny, S. R. Heap, & R. H. Cornett (ASP, San Francisco), p. 273

Gilbank, D. G., Smail, I., Ivison, R. J., & Packham, C. 2003, MNRAS, submitted

Iye, M. et al. 2000, PASJ, 52, 9

Iye, M. et al. 2003, ApJ, 590, 770

Jimenez, R., Padoan, P., Dunlop, J. S., Bowen, D. V., Juvela, M., & Matteucci, F. 2000, ApJ, 532, 152

Katz, N., & Gunn, J. E. 1991, ApJ, 377, 365

Kauffmann, G., & Charlot, S. 1998, MNRAS, 297, 23

Kauffmann, G., & Haehnelt, M. 2000, MNRAS, 311, 576

Knopp, G. P., & Chambers, K. C. 1996, ApJS, 109, 367

Kobayashi, N., et al. 2000, in Proc. SPIE 4008: Optical and IR Telescope Instrumentation and Detectors, ed. M. Iye & A. F. Moorwood, p. 1056

Labbé, I., et al. 2003, ApJ, 591, L95

Liu, M. C., et al. 2000, AJ, 119, 2556

Miyazaki, M. et al. 2003, ApJ, submitted (astro-ph/0210509)

Moriando, G., Cimatti, A., & Daddi, E. 2000, A&A, 364, 26

Motohara, K., et al. 2002, PASJ, 54, 315

Nolan, L. A., Dunlop, J. S., & Jimenez, R. 2001, MNRAS, 323, 385

Nolan, L. A., Dunlop, J. S., Jimenez, R., & Heavens, A. F. 2003, MNRAS, 341, 464

Navarro, J. F., & White, S. D. M. 1994, MNRAS, 267, 401

Peacock, J. A. 1999, Cosmological Physics, (Cambridge U. Press, Cambridge), p. 413

Peebles, P. J. E. 2002, ASP Conf. Ser., 283, 351

Peng, C. Y., Ho, L. C., Impey, C. D., & Rix, H.-W. 2002, AJ, 124, 266

Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525

Sérsic, J.-L. 1968, Atlas de Galaxias Australes, (Observatorio Astronomico, Cordoba)

Smith, G. P., et al. 2002a, MNRAS, 330, 1

Smith, G. P., Smail, I., Kneib, J.-P., Davis, C. J., Takamiya, M., Ebeling, H., & Czoske, O. 2002b, MNRAS, 333, L16

Soifer, et al. 1999, AJ, 118, 2065

Spinrad, H., Dey, A., Stern, D., Dunlop, J, Peacock, J., Jimenez, R., & Windhorst, R. 1997, ApJ, 484, 581

Steinmetz, M., & Navarro, J. F. 2003, New Astron., 8, 557

Stockton, A. 2001, in Astrophysical Ages and Timescales, ASP Conf. Ser., 245, ed. von Hippel et al., (ASP, San Francisco), p. 517

Stockton, A., Kellogg, M., & Ridgway, S.E. 1995, ApJ, 443, L69

Thomas, D., & Kauffmann, G. 1999, in Spectrophotometric Dating of Stars and Galaxies, ASP Conf. Ser., 192, ed. I. Hubeny, S. R. Heap, & R. H. Cornett, p. 261.

Thomas, D., Maraston, C., & Bender, R. 2002, in Reviews in Modern Astronomy, ed. R. E. Schielicke, v. 15, p. 219 [astro-ph/0202166]

Tinsley, B., & Gunn, J. 1976, ApJ, 203, 52

Trager, S. C., Faber, S. M., Worthey, G., & González, J. J. 2000, AJ, 120, 165

Treu, T., Stiavelli, M., Bertin, G., Casertano, S., & Møller, P. 2001, MNRAS, 326, 237

Treu, T., Stiavelli, M., Casertano, S., Møller, P., & Bertin, G. 2002, ApJ, 564, L13

van Dokkum, P. G., & Ellis, R. S. 2003, ApJ, 592, L53

van Dokkum, P. g., Franx, M., Fabricant, D., Illingworth, G. D., & Kelson, D. D. 2000, ApJ, 541, 95

Worthey, G., Faber, S. M., & González, J. J. 1992, ApJ, 398, 69

Yan, L., & Thompson, D. 2003, ApJ, 586, 765